

PILESEPTINE GEL

INTERNAL - GEL

Medical device

INDICATION: Topical dressing for internal hemorrhoids (piles and hemorrhoids located slightly inside the anal orifice).

PRESENTATION: One tube containing 50 ml gel for topical application.

COMPOSITION: Glycerol, Aqua, Pilecyanidin-I (Extracts of Vitis Vinifera and Vaccinium Macrocarpon), Linum Usitatissimum oil, Mentha Piperita essential oil.

MODE OF ACTION:

Pileseptine gel for internal application is a viscous, osmotically active hypertonic solution. When applied on the surface of piles, Pileseptine forms a film over the hemorrhoids and mechanically attracts hypotonic fluid present below the injured surface. This strong outward flow of liquid from the tissue helps to reduce edema and consequently the size of the hemorrhoid. This liquid outflow equally cleans the hemorrhoidal area free of contaminants, and keeps the surface clean and hydrated, which consequently reduces pain, irritation, itching, and burning sensation. Progressive reduction in the size of the hemorrhoid and a clean environment facilitates hemorrhoidal regression.

DIRECTIONS: For topical application only on internal hemorrhoids. After washing your hands, lying on your bed, remove the tip of the tube and press the tube to deliver some product onto your clean, dry finger (1-2 ml). Insert gently the tip of the finger into the rectum up to the lesion and apply the product directly on the lesion. Repeat the same procedure after 2-3 minutes. Wipe off the anal surface with a tissue paper. Recommended posology: 3 to 4 applications per day.

PRECAUTIONS: Wash hands with soap and warm water before and after use. This product is for topical use only for internal hemorrhoids. It should not be swallowed. Do not use PILESEPTINE gel if you are sensitive or allergic to any of the ingredients used to prepare this product. Follow the recommended posology. Seek medical advice if you see no improvement within 7-10 days of use. Keep out of reach of children. Store at room temperature, do not freeze or refrigerate. Do not use the product if the tube is damaged, or after the expiry date.

SIDE EFFECTS: PILESEPTINE gel has no known side effects and is not dangerous if swallowed. In some cases, a slight irritation may be observed which disappears after a few minutes. PILESEPTINE gel can be used in children over 3 years of age, the product being applied by an adult. Safety of use in pregnant or lactating women has not been established. In absence of sufficient data, we do not recommend using this product during pregnancy or lactation.

MANUFACTURER
VITROBIO
ZAC de Lavour
63500 ISSOIRE, France

DISTRIBUTOR



MADE in FRANCE

COMMERCIAL NAMES: PILESEPTINE GEL.

In order to further improve the quality and efficacy of this product, we would greatly appreciate if you could complete and send us, along with your suggestions, the questionnaire on our website www.vitrobio.com
(Questionnaire– Select Language– Piles).
Thank you.